



General *Virucide, Bactericide, Tuberculocide, Fungicide, Sanitizer

When used as directed, this product is highly effective against a wide variety of pathogenic microorganisms including bacteria, antibiotic resistant bacteria, viruses and fungi.

HOSPITAL DISINFECTANT: Bactericidal in the presence of 5% serum load and 1 minute contact time at 20°C (68°F) on hard, non-porous environmental surfaces.

BACTERICIDAL ACTIVITY:

Pseudomonas aeruginosa (ATCC 15442) Escherichia coli 0157:H7 (ATCC 35150) Acinetobacter baumannii (ATCC 19606) Staphylococcus aureus (ATCC 6538) Staphylococcus aureus MRSA (ATCC 33592) Clostridium difficile (vegetative)(ATCC 9689) Salmonella enterica (ATCC 10708) Enterococcus faecalis VRE (ATCC 51575)

*VIRUCIDAL in the presence of 5% serum load and 1 minute contact time at 20°C (68° F) on hard, non-porous environmental surfaces.

Viruses:

Poliovirus Type 1, Strain Brunhilde (ATCC VR-1000)
Human Coronavirus (ATCC VR-740)
Herpes Simplex Virus, Type 2 (HSV-2) (ATCC VR-734)
Hepatitis C Virus (HCV) (acquired from National Veterinary
Services Laboratories)
HIV-1 (AIDS Virus), Strain HTLV-III_B (HIV-1) (acquired from Adv.
Biotechnologies, Inc.)
Influenza A/Hong Kong (ATCC VR-544)
Rhinovirus Type 37, Strain 151-1 (ATCC VR-1147)
Hepatitis B Virus (HBV) (acquired from Hepadnavirus Testing)
Feline Calicivirus, Strain F9 (ATCC VR-782)
Herpes Simplex Virus, Type 1 (HSV-1) (ATCC VR-733)
Rotavirus WA (Aquired from University of Ottawa)

Kills Feline Calicivirus (surrogate for Norovirus) when used as directed on hard, non-porous inanimate surfaces with a 1 minute contact time.

TUBERCULOCIDAL in the presence of a 5% serum load and 5 minute contact time at 20°C (68°F) on hard, non-porous environmental surfaces.

Mycobacterium bovis (Tb) (OT 451C150)

FUNGICIDAL - This product is fungicidal against the pathogenic fungi, *Trichophyton mentagrophytes* (ATCC 9533) (the fungus which causes Athlete's Foot) in the presence of a 5% serum load and 10 minute contact time at 20°C (68°F) when used as directed on hard surfaces found in bathrooms, shower stalls, locker rooms, or other clean, non-porous, hard surfaces commonly contacted by bare feet.

Trichophyton mentagrophytes (ATCC 9533)

Non-Food Contact Surface Sanitizer for hard, non-porous surfaces – in the presence of a 5% serum load and 30 second contact time at 20°C (68°F) on hard, non-porous environmental surfaces.

Klebsiella pneumoniae (ATCC 4352) Staphylococcus aureus (ATCC 6538) Escherichia coli 0157:H7 (ATCC 35150) Salmonella enterica (ATCC 10708) Staphylococcus aureus MRSA (ATCC 33592) Pseudomonas aeruginosa (ATCC 15442) Enterococcus faecalis VRE (ATCC 51575)

Malodor Counteractancy – eliminates odors and odor-causing bacteria in restroom areas, behind and under sinks and counters, storage areas and other places where bacterial growth can cause malodors.

For use as a Non-Food Contact Sanitizer: Pre-clean heavily soiled areas. Apply Solution with cloth, disposable wipe or mop to hard, non-porous environmental surfaces. Allow the surface to remain wet for 30 seconds. Wipe surfaces dry, rinse or allow to air dry. Change solution after each use (application).

EPA Reg. No. 70627-56

©2008 JohnsonDiversey, Inc., Sturtevant, WI 53177-0902 U.S.A.